

**Supplementary File 3. Treatments received for prostate cancer (n=546)**

	<b>Total</b>	<b>Surgery</b>	<b>RT+ADT</b>	<b>Other</b>
Active surveillance*	<b>30 (5%)</b>	15**	2**	13 (8%)
Watchful waiting*	<b>19 (3%)</b>	6**	4**	9 (5%)
Chemotherapy	<b>7 (1%)</b>	n/a	n/a	7 (4%)
Cryotherapy	<b>6 (1%)</b>	n/a	n/a	6 (4%)
EBRT	<b>137 (25%)</b>	n/a	91 (77%)	46 (27%)
HIFU	<b>4 (1%)</b>	n/a	0	4 (2%)
ADT	<b>182 (33%)</b>	n/a	118 (100%)	64 (38%)
Brachytherapy (all)	<b>40 (7%)</b>	n/a	11 (9%)	29 (17%)
<i>Permanent seed</i>	27 (5%)	n/a	5 (4%)	22 (13%)
<i>Temporary</i>	13 (2%)	n/a	6 (5%)	7 (4%)
Radical prostatectomy (all)	<b>310 (57%)</b>	258 (100%)	n/a	52 (31%)
<i>Standard laparoscopic</i>	102 (19%)	92 (36%)	n/a	10 (6%)
<i>Robotic-assisted laparoscopic</i>	73 (13%)	69 (27%)	n/a	4 (2%)
<i>Open</i>	135 (25%)	97 (38%)	n/a	38 (22%)
<b>Total (% of total)</b>	<b>546 (100%)</b>	<b>258 (47%)</b>	<b>118 (22%)</b>	<b>170 (31%)</b>

ADT = androgen deprivation therapy; EBRT = external beam radiotherapy; HIFU = high-intensity focused ultrasound; RT = radiotherapy (including brachytherapy)

Percentages in brackets are a percentage of each column, unless otherwise stated. Respondents could select more than one answer except if brachytherapy (all) or radical prostatectomy (all) was selected, then only one subtype could then be chosen.

"Total" All men with ED after prostate cancer treatment

"Surgery" Men whose only radical treatment was radical prostatectomy

"RT+ADT" Men who only had radiotherapy (EBRT and/or brachytherapy) plus ADT

"Other" All men who received a treatment other than "surgery" alone or "RT + ADT" alone

\* Includes patients who started on active surveillance/watchful waiting who went on to receive radical treatment.

\*\* Data excluded from the percentage calculations.